

Coral Bleaching and Ocean "Hot Spots"

Global sea-surface temperature maps show that mass coral-reef bleaching episodes between 1983 and 1991 followed positive anomalies more than 1°C above long-term monthly averages ("hot spots") during the preceding warm season. Irregular formation, movement, and disappearance of hot spots make their detailed long-term prediction impossible, but they can be tracked in real time from satellite data. Monitoring of ocean hot spots and of coral bleaching is needed if the Framework Convention of Climate Change is to meet its goal of protecting the most temperature-sensitive ecosystems.





Figure 1a. Reef regions of the world, divided into 10 major reef provinces of roughly comparable area. These regions do not necessarily correspond to biogeographic provinces.

REGION	YEAR	79	80	81	82	83	84	85	86	87	88	89	90	91
1. Atlantic-Central		X							X		X	X	X	
2. Atlantic-Peripheral											X		X	X
3. Pacific-NorthWest			X		X									X
4. Pacific-SouthWest				X					X				X	
5. Pacific-East					X									X
6. Pacific-SouthEast						X			X					X
7. Pacific-Central								X	X				X	
8. Indian-West					X			X	X					
9. Indian-Central									X					
10. Indian-East												X	X	
REGION	YEAR	79	80	81	82	83	84	85	86	87	88	89	90	91

Figure 1b. Major bleaching events reported in reef regions since 1979. Bleaching events are taken from published citations and summaries in reference 6, updated by more recent reports from divers and scientists around the world compiled by Ernest Williams (pers. comm.). Local bleaching events are included until 1982, but not afterwards, making any increase in events conservative estimates.


Figure 1c. Frequency of worldwide bleaching events since 1979. In the early 1980s, most reported bleaching events took place in only one reef province per year. In the late 1980s, bleaching events occurred at sites in around half of all reef provinces every year.




CORAL BLEACHING EVENTS

Coral-reef bleaching (1–13), or expulsion of symbiotic zooxanthellae algae, became more frequent, widespread, and severe in the 1980s (6–13). Zooxanthellae provide reef corals most of their carbon (14, 15), limestone depositing ability (16), and color (17). Bleaching causes corals to turn white or pale, because loss of pigment allows the limestone skeleton to become visible through transparent tissues. Local bleaching due to poor water circulation (1–3) or freshwater and sediment from rivers (4,

GLOBAL BLEACHING TRENDS
1979 - 1991







Closeup of a *Montastrea cavernosa* coral head in Negril, Jamaica, in September 1992, slowly regaining pigmentation after being bleached for more than a year. Polyps are around 1 centimeter across. Photo: P.D. Goreau.

5) cannot account for the spatial or temporal pattern of mass bleaching events in the last decade. These events affected shallow and deep corals across large regions, including reefs remote from local stresses (6–13, 18).

Major bleaching events were reported for all major reef provinces in the past decade (Fig. 1a). In 1983, 1987, and 1991 mass bleaching was reported in all tropical oceans (Fig. 1b). El Niño warming events can partially explain this global pattern. Thus, 1983 had the strongest El Niño on record, 1987 had a moderate El

Figure 2. Bleaching events and "hot spot" regions, areas of the ocean with temperature anomalies over 1°C above long-term averages. a) May 1983; b) March 1984; c) September 1986; d) February 1987; e) September 1987; f) August 1988. All hot spots in global monthly average anomaly data in tropical and subtropical zones are shown, but some small hot spots outside these zones are not shown. Major bleaching sites in subsequent months are named and indicated by black circles. Most reports of bleaching at nearby reefs in the same regional hot spots are not shown. Bleaching also took place in the Caribbean in 1989 and 1990 (not shown). It should be noted that although May 1983 water temperatures in the Great Barrier Reef were anomalously high they had already fallen below their maximum seasonal values, and the bleaching event indicated there actually took place in early 1982, before satellite-based global ocean-temperature anomaly maps were available.






Figure 3. Progression of bleaching events and hot spots in 1991. a) February 1991; b) April 1991; c) June 1991; d) August 1991. Mass bleaching events follow the appearance and movement of hot spots as the warmest season of the year progresses from South to North.

Niño, and 1992 had a strong El Niño. However, bleaching took place in both shallow and deep coral areas positively and negatively correlated with El Niño influences (19) and also occurred in non-El Niño years.

In almost all cases, local observers noted that mass bleaching followed extended periods of high temperature, low wind, low cloudiness, and low rainfall. Although high temperatures were thought to be the most probable cause (6–13), rarity of long-term water temperature records in reef habitats made this issue controversial. Disease (20), oxygen toxicity (21), high ultraviolet (UV) light exposure (22), and pollution (23) were also proposed as potential causes. Diseases and parasitic organisms are not seen in histological studies (24, 25). Oxygen toxicity is discounted by experiments showing that zooxanthellae oxygen production ceases at elevated temperatures (26). UV stress is unlikely because bleaching occurs at depths greater than significant UV penetration and because of the presence of UV absorbing pigments in normal and bleached corals (27–29). Experimental data show bleaching is more rapid during high light or UV exposure, but only at

elevated temperature (30–32). Pollution is ruled out because mass bleaching is often more frequent on pristine reefs than inshore and lagoonal reefs subject to freshwater, sewage discharges, and other stresses (6, 10–12, 18, 33). Other potential causes, such as excess turbidity, darkness, or cold, are ruled out by field observations. Nonparametric statistical analysis of reef stress patterns show the spatial distribution of bleaching either does not correlate with or is negatively correlated with all major anthropogenic stresses to reefs (18).

In situ observations and NOAA satellite-derived sea-surface temperature records at seven Atlantic reef locations show that mass bleaching events took place following the warmest periods recorded (34). At each site a threshold monthly mean temperature, approximately 1°C above long-term averages for the warmest months was found, above which bleaching always took place and below which it did not (34). Threshold temperatures are higher at sites with higher mean temperature, indicating that coral populations are adapted to local conditions (34). Mass bleaching was reported in some of the coldest and hottest coral reef areas, showing that both are vulnerable (6). Temperature records at 5 of 7 sites studied in the Caribbean had statistically significant warming trends during the last decade (34). Yet most sites also had significantly decreasing monthly temperature variability and seasonal temperature ranges, with the greatest warming occurring in winter (34).

BLEACHING AND OCEAN HOT SPOTS


We have identified areas where ocean surface temperatures exceeded long-term averages by more than 1°C ("hot spots") during the warmest months preceding major reported bleaching events (Figs 2 and 3). Data are derived from global ocean-temperature anomaly maps published in the NOAA Climate Diagnostics Bulletin (35, 36), and from published bleaching reports, supplemented by field observations and personal communications from divers. Because bleaching rates are much higher at elevated temperatures (30–32), very hot intervals which are too short to appear on monthly mean anomaly data could also cause bleaching, but the fact that most major bleaching events appear to fall within one degree anomalies suggests that such events are fairly rare. This implies that global monthly average sea-surface temperature records are adequate to detect most bleaching conditions, except in the Caribbean, where records with higher spatial resolution are needed (34, 37).

Each region where mass bleaching has been reported since 1983 lay within or on the edge of a warm season hot spot (Figs 2 and 3). Regions where mass bleaching was not reported were not impacted by warm-season anomalies, and hot spots during the cool season were not followed by bleaching. During early 1991, a hot spot formed in the South Pacific outside the coral zone (Fig. 3a). As it disintegrated one portion migrated northwest, passing over the Society Islands, Samoa, and Nauru, before dying out near the equator (Fig. 3b–c). Bleaching and high coral mortality followed in its track (12–13), but were minor in the nearby Tuamotu Islands, which normally have similar climatic conditions to the Society Islands, but lay outside the 1991 hot spot (12). Bleaching later in 1991 at North Pacific, Indian Ocean, and Atlantic sites followed formation of new hot spots (Fig. 3d).

In 1992, in sharp contrast to 1991, very little significant new coral bleaching was reported; 1992 was unusually cold worldwide because high altitude aerosols from the eruption of Mount Pinatubo reflected sunlight back to space. Although El Niño warming affected Eastern Pacific surface waters, the warm-water pool was not suitably timed or located to cause bleaching in major reef areas. Hot spots were virtually absent elsewhere in 1992, but other local stresses to reefs continued to increase. Despite little new bleaching in 1992, the great majority of two of the major reef-building species in Jamaica, *Montastrea cavernosa* and *Siderastrea*



Completely bleached heads of the major Caribbean reef building coral, *Montastrea annularis*, contrast sharply with an unbleached colony of the branching coral *Acropora palmata* (brown with white tips), following hurricane caused bleaching in Jamaica in 1963. Photo: T.F. Goreau.



Normal (dark brown), partially bleached (light brown), and completely bleached (white) plate corals of several *Agaricia* species and an unbleached encrusting coral (*Porites astreoides*, upper right), Jamaica, 1963. These are among the first photographs ever taken of bleaching. Photo: T.F. Goreau.

siderea, still had not fully recovered from the 1991 bleaching event, as late as early 1993. Because these species recovered more rapidly after the three prior mass bleaching events, they may be weakened from repeated stress and could be more strongly affected by subsequent bleaching. After the Pinatubo stratospheric aerosols settle, and the temporary cooling effect wears off, high tropical ocean temperatures could return. The resumption of mass bleaching following the next suitable hot conditions after the Pinatubo cold event would provide strong confirmation that temperature is the major correlative stress to bleaching (47). Divers should therefore remain alert for coral bleaching if warmer conditions return, as even the most pristine reefs could be affected.

TRACKING AND PREDICTION OF HOT SPOTS

Hot spots, like other extreme weather patterns, form, move, and disappear irregularly and unpredictably; their shape, direction of movement, and duration are extremely variable (38). Chaos in the climate system makes detailed long-term prediction of ocean hot spots unlikely. Extreme atmospheric temperatures usually last only a few weeks, but ocean hot spots typically last a few months due to the greater "memory" of the ocean for temperature changes (39). As a month or more of water temperatures 1°C above average in the warmest months precedes most recorded mass bleaching events, tracking ocean thermal anomalies in real time could allow warning of sites where mass bleaching may be about to start.

There appear no clear latitudinal, regional, or seasonal trends in hot-spot distribution. Model-based projections suggest global warming rates should be greater in polar than equatorial regions, causing assessments of the potential ecological impacts of climate change to focus primarily on colder regions (40). Many tropical organisms, including both corals and major tropical plant crops, show strong stress responses under high temperature conditions, even when all other environmental factors are adequate. Neglecting ecophysiological impacts of high temperature on tropical biomes is especially unwise because most tropical organisms live much closer to their upper temperature limits than those in cold zones, and so are much more sensitive to small temperature increases (41).

Widespread evidence that coral reefs around the world underwent episodic high temperature stress over the past decade makes bleaching a harbinger of reef ecosystem stress from ocean warming (6–10, 42). Global warming rates over the past century and a half have been quite similar over oceans and land, and Northern and Southern Hemispheres, with the 1980s marking the highest values in the record (43). Previously unaffected regions could be impacted if hot spots continue with the frequency of the 1980s, and the health of coral-reef ecosystems globally could be compromised if hot spots become more frequent and intense, as predicted by model global warming scenarios (38). Bleached corals either fail to grow and reproduce until after they recover during the following cool season (44), or die if stress is excessive (1, 12, 13, 30–32). Continued or increased mass coral bleaching would seriously impact marine biodiversity, fisheries, tourism, shore protection, and the ability to adapt to rising sea level in the over 100 countries where coral reefs are major natural and economic resources.

CORAL REEFS AND THE CLIMATE CHANGE CONVENTION

Coral reefs are perhaps the most vulnerable ecosystems to rising temperatures, sea levels, soil erosion, and to excess nutrients from sewage and fertilizers. This gives them a central role if the Climate Change Convention, signed in Rio de Janeiro, is to meet its own stated goal of protecting Earth's most climatically sensitive ecosystems. However, the Convention failed to require immediate monitoring of the most sensitive ecosystems, or to set acceptable limits of climate change at values below their tolerance limits. Global monitoring of both hot spots and coral bleaching should be promptly begun to determine the current extent and impact of large-scale temperature stress and whether current rates of climate change are within the ability limits of reef corals to adapt. If not, more stringent limits on greenhouse-gas emissions may be required.

If global warming continues almost all ecosystems can be replaced by migration of species from lower latitudes, except for the warmest ecosystems. These have no source of immigrants already adapted to warmer conditions. Their species must evolve new environmental tolerances if their descendants are to survive,

a much slower process than migration. Research on physiological tolerance and adaptation capability of tropical organisms should be central to any effort to protect the most biodiverse, productive, and beautiful tropical ecosystems; coral reefs and rainforests. Large-scale funding for long-term research in these habitats is critically needed yet woefully inadequate in current international environmental protection and development policies.

References and Notes

1. Mayor, A.G. 1918. Toxic effects due to high temperature. *Carnegie Inst. Wash. Pap. Mar. Biol.* 12, 175–178.
2. Yonge, C.M. and Nicholls, A.G. 1931. Studies on the physiology of corals. *Great Barrier Reef Exp. Sci. Pap.* 1, 135–211.
3. Jaap, W. 1979. Observation on zooxanthellae expulsion at Middle Sambo Reef, Florida Keys. *Bull. Mar. Sci.* 29, 414–422.
4. Goreau, T.F. 1964. Mass expulsion of zooxanthellae from Jamaican reef communities after Hurricane Flora. *Science* 145, 383–386.
5. Goenaga, C. and Canals, M. 1979. Relacion de mortandad masiva de *Millepora complanata* (Cnidaria, Anthozoa) can alta pluviosidad y escorrienta del Rio Fajardo en Cayo Ahogado. *Symp. Nat. Res. San Juan Puerto Rico* 6, 84–95. (In Spanish).
6. Williams, E.H. and Bunkley-Williams, L. 1990. The world-wide coral bleaching cycle and related sources of coral mortality. *Atoll Res. Bull.* 335, 1–71. See also the many additional published and unpublished bleaching reports cited in references 6 and 34 which can not be repeated here for reasons of space.
7. Jokiel, P.L. and Coles, S.L. 1990. Response of Hawaiian and other Indo-Pacific corals to elevated temperatures. *Coral Reefs* 8, 155–162.
8. Glynn, P.W. 1991. Coral reef bleaching in the 1980s and possible connections with global warming. *Trends Ecol. Evol.* 6, 175–179.
9. Hayes, R.L. and Goreau, T.J. 1991. Tropical coral reef ecosystem as a harbinger of global warming. *World Resour. Rev.* 3, 306–322.
10. Goreau, T.J. 1990. Coral bleaching in Jamaica. *Nature* 343, 417.
11. Goenaga, C. and Canals, M. 1990. Island-wide coral bleaching in Puerto Rico. *Carib. J. Sci.* 26, 171–175.
12. Salvat, B. 1992. Blanchissement et mortalité des scleractiniaires sur les récifs de Moorea (archipel de la Société) en 1991. *C. R. Acad. Sci. Paris Ser. II* 314, 105–111.
13. Salvat, B. 1992. The 1991 bleaching event in the Society Islands, French Polynesia. *Proc. 7th Int. Coral Reef Symp.* Guam.
14. Goreau, T.F. and Goreau, N.I. 1960. Distribution of labelled carbon in reef building corals with and without zooxanthellae. *Science* 131, 668–669.
15. Muscatine, L. 1973. Nutrition of corals. In: *Biology and Geology of Coral Reefs Vol. 2*. Jones, O.A. and Endean, R. (eds). Academic Press, New York, p. 77–115.
16. Goreau, T.F. and Goreau, N.I. 1959. The physiology of skeleton formation in corals. II. Calcium deposition by hermatypic corals under various conditions in the reef. *Biol. Bull.* 117, 239–250.
17. Kleppel, G.S., Dodge, R.E. and Reese, C.J. 1989. Changes in pigmentation associated with the bleaching of stony corals. *Limnol. Oceanogr.* 34, 1331–1335.
18. Goreau, T.J. 1992. Bleaching and reef community change in Jamaica: 1951–1991. *Am. Zool.* 32, 683–695.
19. Trenberth, K. 1989. TOGA and atmospheric processes. In: *Understanding Climate Change*. Berger, A., Dickinson, R.E. and Kidson, J.W. (eds). American Geophysical Union, Washington DC, p. 117–125.
20. Williams, E., Goenaga, C. and Vicente, V. 1987. Mass bleaching on Atlantic coral reefs. *Science* 238, 877–878.
21. Sandeman, I. 1988. Zooxanthellae physiology and a suggested irradiance temperature mechanism for bleaching. *Proc. Assoc. Island Mar. Labs. Carib.* 21, 50.
22. Lesser, M.P., Stochaj, W.R., Tapley, D.W. and Shick, J.M. 1990. Bleaching in coral reef anthozoans: effects of irradiance, ultraviolet radiation, and temperature on the activities of protective enzymes against active oxygen. *Coral Reefs* 8, 225–232.
23. D'Elia, C., Buddemeier, R. and Smith, S. 1991. *Workshop on Coral Bleaching, Coral Reef Ecosystems, and Global Change: Report of Proceedings*. Maryland Sea Grant College, College Park, Maryland.
24. Glynn, P.W., Peters, E.C. and Muscatine, L. 1985. Coral tissue microstructure and necrosis: relation to catastrophic coral mortality in Panama. *Dis. Aquat. Organisms* 1, 29–37.
25. Hayes, R.L. and Bush, P. 1990. Microscopic observations of recovery in the reef building scleractinian coral *Montastrea annularis*, after bleaching on a Cayman reef. *Coral Reefs* 5, 201–204.
26. Iglesias-Prieto, R., Matta, J., Robins, W. and Trench, R.K. 1993. The photosynthetic response to elevated temperature in the symbiotic dinoflagellate *Symbiodinium microadriaticum* in culture. *Proc. Nat. Acad. Sci.* 89, 10302–10305.
27. Jokiel, P. and York, R.H. 1984. Solar ultraviolet photobiology of the reef coral *Pocillopora damicornis* and symbiotic zooxanthellae. *Bull. Mar. Sci.* 32, 301–305.
28. Dunlap, W.C., Chalker, B.E. and Oliver, J.K. 1986. Bathymetric adaptations of reef-building corals at Davies Reef, Great Barrier Reef, Australia. III. UV-B absorbing compounds. *J. Exp. Mar. Biol. Ecol.* 104, 239–248.
29. Siebeck, O. 1988. Experimental investigation of UV tolerance in hermatypic corals (Scleractinia). *Mar. Biol. Prog. Ser.* 43, 95–103.
30. Jokiel, P. and Coles, S. 1977. Effect of temperature on the mortality and growth of Hawaiian reef corals. *Mar. Biol.* 43, 201–208.
31. Hoegh-Guldberg, O. and Smith, G.J. 1989. Light, salinity, and temperature and the population density, metabolism, and export of zooxanthellae from *Stylophora pistillata* and *Seriatopora hystrix*. *J. Exp. Mar. Biol. Ecol.* 129, 279–303.
32. Glynn, P.W., Imai, R., Sakai, K., Nakano, Y. and Yamazoto, K. 1992. Experimental response of Okinawan (Ryukyu Islands, Japan) reef corals to high sea temperature and UV radiation. *Proc. 7th Int. Coral Reef Symposium*, Guam.
33. Goreau, T.J. 1991. Coral reef health in the Negril area. In: *Protecting Jamaica's Coral Reefs*. Reef Relief and Negril Coral Reef Preservation Society, Key West, Florida, p. 30–70.
34. Goreau, T.J., Hayes, R.L., Clark, J.W., Basta, D.J. and Robertson, C.N. 1993. Elevated sea surface temperatures correlate with Caribbean coral reef bleaching. In: *A Global Warming Forum: Scientific, Economic and Legal Overview*. Geyer, R.A. (ed.). CRC Press, Boca Raton, Florida, p. 225–255.
35. *Climate Diagnostics Bulletin 1983–1992*. US Dept. of Commerce, Washington DC.
36. Methods: Hot spot maps were prepared by tracing outlines of the 1°C anomalies from global monthly average maps for the hottest month preceding major reported bleaching events. NOAA Oceanographic Monthly Summary temperature data (reference 35) are based on blended satellite and shipboard raw data, processed by spatial and temporal averaging over several degrees of latitude and longitude (R. Reynolds, pers. comm.), and the anomaly determined by comparison with long term oceanographic data sets (Reynolds, R. 1988, A real-time global sea surface temperature analysis. *J. Climate* 1, 75–86). Satellite-derived sea surface temperature data sets have been controversial (Strong, A.E. 1989. Greater global warming revealed by satellite-derived sea surface temperature trends. *Nature* 338, 642–645) because signals in the thermal infrared channel of satellite radiometers require correction for interference by clouds and stratospheric aerosols caused by volcanic eruptions. These effects cause satellite data to somewhat underestimate actual surface temperatures, and corrections for them have relatively minor effects on reported trends (Reynolds, R.W., Folland, C.K. and Parker, D.E. 1989. Biases in satellite-derived sea surface temperature data. *Nature* 341, 728–731; Robock, A. and Strong, A.E. 1989. Satellite data contamination. *Nature* 341, 695).
37. Although global thermal anomaly maps are successful at identifying hot spots preceding mass bleaching in the Pacific and Indian Oceans they do not work well in the Caribbean region (marked by a rectangle in Figures 2 and 3). The Caribbean is apparently a special case because the effective pixel size of the global maps is actually several hundred km. (R. Reynolds, pers. comm.), which is too large to resolve the Caribbean's complex fine-scale current flows and temperature patterns (A. Strong, pers. comm.). High resolution Caribbean satellite ocean temperature images with spatial resolution of 4–50 km are also available which show hot spots that are not visible in lower resolution data (Strong, A.E. and McCormack, R.C. 1991. Coral bleaching and sea surface temperatures. *Proc. Oceanog. Soc. Conf.*, St. Petersburg, Florida). An intermediate resolution satellite data set, the Ocean Features Analysis, prepared by NOAA for identifying eddies in the Gulf Stream region (J. Clark, pers. comm.) was directly calibrated against measured *in situ* surface water temperatures in Jamaica and Grand Cayman, and found generally accurate to around 0.2°C (34). This data set also resolves Caribbean hot spots which do not show up in the low resolution Oceanographic Monthly Summary data. Low resolution Caribbean data (Atwood, D.K., Sylvester, J.C., Corredor, J.E., Morell, J.M., Mendez, A., Nodal, W.J., Huss, B.E. and Foltz, C. 1988. Sea surface temperature anomalies for the Caribbean, Gulf of Mexico, Florida reef track, and Bahamas considered in light of the 1987 regional coral bleaching event. *Proc. Ass. Island Mar. Labs. Carib.* 21, 47) are not well correlated with *in situ* surface water temperature measurements, and underestimate observed values by up to several degrees (34). Confusion between these data bases has resulted in publication of high resolution Caribbean temperature images which accurately show hot spots above 31°C at bleaching sites (see images in Brown, B., Ogden, J. 1993. Coral bleaching. *Sci. Am.* 268, 64–70 which were provided by A.E. Strong), juxtaposed with conclusions based on low resolution maps that show spuriously low temperatures around 29 Celsius at the same time, which were used to conclude that bleaching is not linked to high temperature (Atwood, D., Hendee, J. and Mendez, A. 1992. An assessment of global warming stress on Caribbean coral reef ecosystems. *Bull. Mar. Sci.* 51, 118–130).
38. Hansen, J., Fung, I., Lacis, A., Rind, D., Lebedeff, S., Ruedy, R. and Russell, G. 1988. Global climate changes as forecast by Goddard Institute for Space Studies three-dimensional model. *J. Geophys. Res.* 93, 9341–9346.
39. Sciremamano, F. 1979. A suggestion for the presentation of correlations and their significance levels. *J. Phys. Oceanogr.* 9, 1273–1276.
40. Houghton, J.T., Jenkins, C.J. and Ephraums, J.J. 1990. *Climate Change: the IPCC Assessment*. Cambridge U. Press, Cambridge, UK.
41. Emiliani, C.E., Kraus, E.B. and Shoemaker, E.M. 1981. Sudden death at the end of the Mesozoic. *Earth Planet. Sci. Lett.* 55, 317–334.
42. Glynn, P.W. and Weerdt, W.H. 1991. Elimination of two reef-building hydrocorals following the 1982–83 El Niño warming event. *Science* 253, 69–71.
43. Houghton, J.T., Callander, B.A. and Varney, S.K. 1992. *Climate Change 1992: The Supplementary Report to the IPCC Scientific Assessment*. Cambridge U. Press, Cambridge, UK.
44. Goreau, T.J. and Macfarlane, A.H. 1990. Reduced growth rate of *Montastrea annularis* following the 1987–1988 coral bleaching event. *Coral Reefs* 8, 211–216.
45. Acknowledgements: We gratefully thank Dr. Ernest Williams of the University of Puerto Rico at Mayaguez for discussion and information from his compilation of diver reports of bleaching and major marine environment disturbances. We also thank A. Strong, R. Reynolds, J. Clark, R.K. Trench and N.I. Goreau for useful discussions, and the reviewers for helpful comments.
46. This paper is dedicated to the memory of Dr. Carlos Goenaga, a brilliant researcher on corals and bleaching at the University of Puerto Rico at Mayaguez, tragically killed by a wave while working on March 10, 1993.
47. Mild bleaching took place in 1993 in Jamaica, Florida, and the Bahamas following at small Western Caribbean hot spot. The first hot spot in the Pacific reached the Society Islands in February 1994, and mass bleaching began in March.
48. First submitted 8 March, 1993, accepted for publication after revision 12 July, 1993.

Thomas J. Goreau, PhD is president of the Global Coral Reef Alliance and Science Advisor to the Negril Coral Reef Preservation Society. He works primarily on long-term changes in coral reefs in Jamaica and policies for reef protection. His address: Global Coral Reef Alliance, 324 North Bedford Road, Chappaqua, N.Y. 10514 USA. Raymond L. Hayes, PhD is Chairman of Anatomy at Howard University Medical School and former Chairman of Marine Sciences at the University of the Virgin Islands. He works mainly on molecular, biochemical, and cellular responses of corals. They have done field studies and experimental research on coral bleaching in the Caribbean and in the South Pacific. His address: Department of Anatomy, College of Medicine, Howard University, Washington DC 20059, USA.